

# Impacts on Mangrove Distributions in the Firth, and Implications

Carolyn Lundquist and Sanjay Wadhwa, NIWA, Hamilton


(presented by Graham McBride, NIWA, Hamilton)



# What are projected changes in mangrove distributions in the Firth of Thames?


- Climate Change likely to result in changes in:
  - Sedimentation rates
  - Sea level rise
  - Wave/storm frequency
  - Wind direction and strength
- Key driver of change in mangrove distribution is change in bed level height
- What is the likely impact on distributions of coastal habitats (mangroves, salt marsh) of these combined influences?

Study area: Firth of Thames/Waihou R/Gulf





# NZ Mangroves

- Mangrove forests (*Avicennia marina*) occur in upper North Island estuaries
- **Modern forests** developed over last ~150 yr = period of increased catchment erosion & accelerated estuary infilling
- **Mangrove-habitat expansion** averaging  $4\% \text{ yr}^{-1}$  (last ~70 yr ) = **displaced** other habitats
- **Intertidal zone** above mean sea level (**MSL**) = physiological tolerance to immersion
- **Must maintain position relative to MSL** as sea-level rises (surface elevation gain)




# Key stressors: impacts, tipping points and implications for policy and management

- Climate changes
  - Temperature
  - **Sea Level Rise**
  - **Storm/Wave/Wind Climate**
  - **Rainfall**
  - **Catchment Sedimentation**



- Changes in biological habitat distribution (and Ecosystem Services provided by them)



- Changes in bed level height through sediment deposition and/or erosion

# Model scenarios


- 3D Deltares model
  - Water Column sediment deposition model
- 

Projected changes in high tide water depth for:

- 3 'positions'
  - intertidal wetting and drying;
  - edge of intertidal zone;
  - subtidal platform
- 4 RCPs ( $W/m^2$ )
  - 2.6
  - 4.5
  - 6.0
  - 8.5
- 3 sedimentation rates
  - 300  $g/m^3$  (in deep water)
  - 20% increase
  - 20% decrease


# Convert projected changes in water depth to mangrove distribution

Projected changes in high tide water depth




- LiDAR to determine area in each depth band for 25cm increments from 3m above to 3m below '0 m' water depth
- Convert model projections of high tide water depth to LiDAR 'depths'
- Determine depth bands relative to mangrove ecological values;
  - Saltmarsh band
  - High suitability for mangroves (~80% in current FoT mangrove distribution)
  - Low suitability (above mean water depth, but typically low colonisation success by mangroves) (~10% colonised in current FoT mangrove distribution, though biology suggests these are potentially suitable for mangroves)
  - Subtidal - unsuitable
- Calculate revised depth bands and area covered after conversion using projected changes in high tide water depth






# Example: change in mangrove distribution






# Interactions between sedimentation and sea level rise


RCP 6.0, no change sedimentation



RCP 6.0, 20% increase sedimentation



RCP 6.0, 20% decrease sedimentation



# Summary of results

- Sedimentation and sea level rise interact to determine changes in coastal habitat distributions under different climate change scenarios
- In some scenarios, sedimentation is not sufficient to maintain mangrove distributions
- In other scenarios, increased sedimentation builds up bed level height resulting in increased mangrove distribution, likely at expense of saltmarsh and shallow subtidal habitats



# Implications

- Landuse policies and practice need to be cognisant of the *primary role played by sediment runoff* on the evolution of bathymetry and water depth in the Firth over the next 100 years.
  - In contrast to the Waihou River study where SLR is rather dominant
  - SLR and sedimentation show signs of being in a state of equilibrium (differences between RCPs generally small)
- Nonetheless, water levels will rise of course, so waters' edge activities and infrastructure could be impaired (roads, railways,...).
- Further work needed to:
  - analyse effect of various land use policies
  - project changes in fisheries distribution, spawning, land flooding, storm surges

# Next steps

- Sedimentation is important to maintain mangroves with respect to sea level rise.
- Barriers to seaward migration?
- Impacts on ecosystem services provided by coastal habitats?

